New Signals and Backgrounds in Direct Detection Experiments

N3AS Seminar
July 12, 2022

Mukul Sholapurkar
UC San Diego

Introduction

- Strong evidence for the existence of dark matter through its gravitational interactions
- The big question: does dark matter interact with SM particles through forces other than gravity?
- Several experiments have been probing various models of dark matter in different ways

Dark Matter (DM) Detection Strategies



Dark Matter (DM) Detection Strategies



Channels of Direct Detection

Recent Detection Method (probe sub-GeV DM)

Proposed Detection Method (probe sub-MeV DM)

Backgrounds?



Traditional Detection Method (~ GeV DM mass and above)

Well-Understood Radiogenic and Cosmogenic Backgrounds above ~ keV recoils

*arXiv:1910.08092 Trickle, Zhang, Zurek, Inzani, Griffin

Theory Motivation



Well-motivated targets in sub-GeV parameter space

Sensitivities could be limited by backgrounds

Excesses in Low-Threshold Dark Matter Searches

- Several experiments observe a sizable number of low-energy events
- For example, searches for electron recoils like SENSEI:



*SENSEI, 2020

SENSEI
(Si, Skipper-CCD)

Hypothesis

*arXiv:2011.13939 Du, Egana-Ugrinovic, Essig, **MS**

- High energy particles can interact to create low-energy photons
- These photons can be absorbed to produce low-energy electronic recoil events



- **Cherenkov Radiation**
- Transition Radiation

Concrete Example: SENSEI at MINOS

SENSEI overview:

- Uses silicon Skipper-CCDs to probe sub-GeV DM by precisely measuring ionization





**SENSEI, 2020*

SENSEI at MINOS

SENSEI data:

- **Excellent spatial resolution**
- Can place cuts based on the position of events relative to the positions of high-energy tracks
- Observed ~ 450 1-e events per (gram*day) after applying a 60-pixel (~900 μm) halo-mask cut



*SENSEI, 2020



Radiative Processes Simulation for SENSEI

*paper in prep: Du, Egana-Ugrinovic, Essig, **MS**



Radiative Processes Simulation for SENSEI

*paper in prep: Du, Egana-Ugrinovic, Essig, **MS**

Simulated Tracks



SENSEI data

Radiative Processes Simulation for SENSEI

*paper in prep: Du, Egana-Ugrinovic, Essig, **MS**

Simulated Tracks + Cherenkov



SENSEI data

Radiative backgrounds in SENSEI

*paper in prep: Du, Egana-Ugrinovic, Essig, **MS**

Results:

- SENSEI rate after a 60-pixel halo-mask cut: **450 ± 45 / (gram*day)**
- Estimated Cherenkov contribution: **150 ± 40 / (gram*day)** (including systematics)



Channels of Direct Detection

Recent Detection Method (probe sub-GeV DM)

Proposed Detection Method (probe sub-MeV DM)

Backgrounds?



Traditional Detection Method (~ GeV DM mass and above)

Well-Understood Radiogenic and Cosmogenic Backgrounds above ~ keV recoils

*arXiv:1910.08092 Trickle, Zhang, Zurek, Inzani, Griffin

Theory Motivation



Well-motivated targets in sub-MeV parameter space

Sensitivity projections assuming **background-free** kg-year exposure

Important to estimate backgrounds in these future detectors

*arXiv:2108.03239 Kahn, Lin

Photon Background

*arXiv:2112.09702 Berghaus, Essig, Hochberg, Shoji, **MS**



Phonons from Coherent Atomic Scattering

$$\frac{d\sigma}{d\Omega d\omega}(q, E_\gamma, \omega) = \frac{d\sigma}{d\Omega}(q, E_\gamma) S(q, \omega)$$

Coherent Atomic Scattering
Cross Section

Dominated by Thompson Cross section
with individual electrons

$$\frac{d\sigma_T}{d\Omega}(q, \theta) \simeq \frac{\alpha^2}{2m_e^2}\left(1 + \cos^2\theta\right)|g(q)|^2$$

*H. Schober 2014

Structure Function

$$S(\vec{q}, \omega) = \sum_f |\langle f|e^{i\vec{q}\cdot\vec{r_N}}|i\rangle|^2 \delta(E_i - E_f - \omega)$$

Can be evaluated in terms of the
phonon density of states of
the material

Phonons from Coherent Atomic Scattering

$$\frac{d\sigma}{d\Omega d\omega}(q, E_\gamma, \omega) = \frac{d\sigma}{d\Omega}(q, E_\gamma) S(q, \omega)$$

Coherent Atomic Scattering
Cross Section

Dominated by Thompson Cross section
with individual electrons

$$\frac{d\sigma_T}{d\Omega}(q, \theta) \simeq \frac{\alpha^2}{2m_e^2}\left(1 + \cos^2\theta\right)|g(q)|^2$$

*H. Schober 2014

Structure Function

$$S(\vec{q}, \omega) = \sum_f \left|\langle f | e^{i\vec{q}\cdot\vec{r_N}} | i \rangle\right|^2 \delta(E_i - E_f - \omega)$$

Can be evaluated in terms of the
phonon density of states of
the material

Final Rate: $$\frac{dR}{d\omega}(\omega) = N_T \sum_i \int d\Omega \frac{d\sigma}{d\Omega d\omega}(q, E_{\gamma_i}, \omega) n_{\gamma_i} v$$

Phonons from Coherent Atomic Scattering



Phonon rates assuming
a typical photon background measured
in a well-shielded
EDELWEISS detector

Could be as high as
~ 100 events per kg.year

Background Conclusions

- Important to identify, characterize and mitigate new backgrounds in low threshold experiments
- Current low-threshold electron recoil experiments could be limited by backgrounds from radiative processes like Cherenkov
- Future experiments that will look for phonons may be limited by backgrounds created by high energy photons
- Mitigation strategies: Increase passive shielding (less high energy photons, muons etc.), Active shielding (correlated high energy events)

Solar Neutrinos as a Signal in Low Threshold Direct Detection Experiments

*arXiv:1801.10159 Essig, **MS**, Yu

Signals from Solar Neutrinos







Need at least a few kg.years of clean exposure in semiconductors

Solar Neutrino Fluxes



Possibility of detecting lower energy tails of solar neutrinos

Solar Metallicity Problem

Combined with Borexino measurement, could give a direct measurement of electron neutrino survival probability

Thanks!